



Can synchronous multisensory looming stimuli bias attentional weights?

Huygelier, H. ¹✉, van Ee, R. ^{1, 2}, Wagemans, J. ¹, Gillebert, C. R. ^{1, 3}

¹ Laboratory of Experimental Psychology, Brain & Cognition, KU Leuven, Belgium

² Philips Research Laboratories, Department of Brain, Body & Behavior, Eindhoven, The Netherlands

³ Department of Experimental Psychology, University of Oxford, United Kingdom



✉ hanne.huygelier@kuleuven.be

@neuropsychylab

www.neuropsychologylab.be

Summary

Background

- Attentional weights are computed which guide selective attention (Bisley & Goldberg, 2010).
- Spatially linked audiovisual information is prioritized (Talsma, Senkowski, Soto-Faraco, & Woldorff, 2010).
- Multisensory processing is more dominant in elderly (Diaconescu, Hasher, & McIntosh, 2013).

Research Question


Can synchronous multisensory stimuli affect attentional weights in a bottom-up fashion?

Conclusion


- Grouping by visual synchronicity aids divided attention.
- Healthy observers do not automatically assign higher attentional weights to audiovisual synchronous stimuli compared to audiovisual asynchronous stimuli.

Stimuli

- Looming frequencies: fast (1.25Hz), slow (0.75Hz)




Block Design



Procedure

Main Experiment



- Task: contrast change detection
- 12 blocks: 44 target trials + 4 catch trials

Synchronicity Discrimination

- 2 looming disks presented (8s): 1 in sync with auditory looming
- Forced choice: which disk in sync with auditory looming?

Participants

Young group

- N = 18 (n = 16)
- Age: Median = 22 years (SD = 3.0)

Older group

- N = 28 (n = 20)
- Age: Median = 70 years (SD = 6.7)
- Neurologically healthy

Synchronicity Discrimination

- High accuracy:
 - Young group: M = 99% (SD = 2.5)
 - Old group: M = 96% (SD = 9.0)

Results main experiment

